

A more detailed experience with the measurement of specific urinary marker proteins has recently been published [4].

AXEL REGENITER, WERNER H. SIEDE, ANDRÉ SCHOLER,
PETER HUBER, NIKOLAUS FRISCHMUTH, and JÜRGEN U. STEIGER
Basel, Switzerland, and Lemgo, Germany

Correspondence to Axel Regeniter, M.D., Kantonsspital Basel, University Hospital, Central Laboratory, Petersgraben 4, CH 4031 Basel, Switzerland.

E-mail: regenitera@uhbs.ch

REFERENCES

1. BAZZI C, PETRINI C, RIZZA V, *et al*: A modern approach to selectivity of proteinuria and tubulointerstitial damage in nephrotic syndrome. *Kidney Int* 58:1732–1741, 2000
2. STEINHAUSLIN F, WAUTERS JP: Quantitation of proteinuria in kidney transplant patients: Accuracy of the urinary protein/creatinine ratio. *Clin Nephrol* 43:110–115, 1995
3. GUDER WG, HOFMANN W: Differentiation of proteinuria and hematuria by single protein analysis in urine. *Clin Biochem* 26:277–282, 1993
4. REGENITER A, SIEDE WH, SCHOLER A, *et al*: Interpreting complex urinary patterns with MDI LABLINK: A statistical evaluation. *Clin Chim Acta* 297:261–273, 2000

Reply from the authors

This letter from Regeniter *et al* about the best method to correctly assess the selectivity of proteinuria is rather confusing. In one instance, they compare the selectivity index with the IgG/transferrin ratio, while in Table 1 and another text mention the selectivity index is compared with the transferrin/IgG ratio. Moreover, the authors do not specify whether the patients they studied had nephrotic proteinuria. The very high rates of unselective proteinuria reported in Table 1, very different from the rates usually found in patients with nephrotic syndrome from primary glomerulonephritis, suggest that most patients tested had low levels of proteinuria. It is well known that in this case the value of the selectivity index is not reliable.

In our study [1] we measured IgG and transferrin by immunonephelometry in second-morning urine samples (**Methods** section) and calculated the selectivity index according to the formula of Cameron and Blandford [2] in 89 patients with nephrotic syndrome. The simple calculation of the IgG/transferrin ratio in the same 89 patients shows values very different from that obtained by the usual formula (Table 1). Moreover, our classification of the selectivity index in patients with nephrotic syndrome and baseline normal renal function fits very well with the functional outcome: remission, high selectivity (HS) 100%; moderate selectivity (MS) 50%; and nonselectivity (NS) 29%; $P = 0.0001$ and progression to

Table 1. Glomerular selectivity assessed by selectivity index vs. urinary IgG/transferrin ratio

	N	SI	IgG/transferrin
SI ≤ 0.10 (HS)	15	0.07 ± 0.02	0.24 ± 0.11
SI $\geq 0.11 \leq 0.20$ (MS)	34	0.15 ± 0.03	0.50 ± 0.25
SI ≥ 0.21 (NS)	40	0.33 ± 0.11	1.17 ± 1.19

Abbreviations are: HS, high selectivity; MS, moderate selectivity; NS, nonselectivity; SI, selectivity index according to the formula of Cameron and Blandford [2]; IgG/transferrin, ratio of IgG over transferrin measured in second morning urine samples.

chronic renal failure, HS 0%, MS 25%, and NS 35%; $P = 0.05$. The selectivity index calculated as urinary IgG/transferrin ratio does not show this type of association. Furthermore, our results in patients with focal segmental glomerulosclerosis and minimal change disease are similar to the results of Laurent *et al* [3], who found that 100% of patients with a selectivity index ≤ 0.07 were steroid-responsive, while all patients with a selectivity index ≥ 0.17 were steroid-resistant. Finally, if, according to the suggestion of Regeniter *et al*, urinary α_2 -macroglobulin is measured, they are measuring three proteins, while we are measuring four proteins (serum and urinary IgG and transferrin). That is not a very big and “laborious” difference!

In conclusion, in our opinion, the selectivity index is clinically useful in patients with nephrotic syndrome and the correct method to assess the selectivity of proteinuria is by using the formula of Cameron and Blandford [2].

Another method used to evaluate the characteristics of both the glomerular and tubular components of proteinuria is to measure, in second morning urine samples, some proteins with different molecular weight [for example, IgG 150 kD and α_1 -microglobulin (α_1 m) 31.8 kD] expressed in milligrams per gram of urinary creatinine (U_{Cr}). In patients with membranous nephropathy (abstract; Bazzi *et al*, *J Am Soc Nephrol* 9, 84A, 1998, updated and submitted for publication) we found that the patients with IgG excretion $< vs \geq 110$ mg/g U_{Cr} have 100% and 20% remission ($P = 0.0001$), respectively, and the patients with α_1 m excretion $< vs \geq 33.5$ mg/g U_{Cr} have 0% vs. 58% progression to chronic renal failure ($P = 0.0001$), respectively.

This is the most simple method to evaluate the quality of proteinuria because only two urinary proteins are measured and their levels that, respectively, reflect the alteration of size-selectivity and the impairment of tubular reabsorption of microproteins, have a high predictive value of functional outcome, at least in patients with idiopathic membranous nephropathy.

CLAUDIO BAZZI, CONCETTA PETRINI, VIRGINIA RIZZA,
GIROLAMO ARRIGO, and GIUSEPPE D'AMICO
Milan, Italy

Correspondence to Claudio Bazzi, M.D., Division of Nephrology and Dialysis, San Carlo Borromeo Hospital, Via Pio II, n. 3, 20153, Milan, Italy.
E-mail: giuseppe.damico@oscb.sined.net

REFERENCES

1. BAZZI C, PETRINI C, RIZZA V, *et al*: A modern approach to selectivity of proteinuria and tubulointerstitial damage in nephrotic syndrome. *Kidney Int* 58:1732–1741, 2000
2. CAMERON JS, BLANDFORD G: The simple assessment of selectivity in heavy proteinuria. *Lancet* 2:242–244, 1966
3. LAURENT L, PHILIPPON C, LAGRUE G, *et al*: Proteinuria selectivity index-prognostic value in lipid nephrosis and related diseases. *Nephron* 65:185–189, 1993

Cited “validation” references for the SphygmoCor device

To the Editor: Covic *et al* describe attempts to measure “central arterial pressure waveforms” in hemodialysis patients in Romania [1]. They claim their noninvasive assessments of “aortic” blood pressure (BP) waveforms using the SphygmoCor device (PWV Medical, Inc., Sydney, Australia) have been validated. However, they cite no data to support this assertion.

The authors write, “The software analytical program also derived in real time from the measured radial artery waveform an aortic BP waveform using a validated transfer function algorithm” [1, p 2636]. However, this is not correct. The two references cited (23 and 24) have nothing to do with the SphygmoCor and have not validated the SphygmoCor’s generalized transfer function (GTF) algorithm.

Reference 23 is a 2-page short report from a 1992 Supplement, a year before both the SphygmoCor radial artery GTF was published [2], and the technique’s United States Patent was granted [3]. Reference 24 did not use the SphygmoCor, but rather involved another GTF developed using a completely different computational technique. This approach has subsequently been shown to be ineffective in 67% of cases, when calibrated noninvasively [4].

A search on Medline reveals a paucity of validation work with the SphygmoCor reported in the literature. Furthermore, no evidence has been provided to support the use of the device in patients with renal failure, let alone in those following hemodialysis. Given this, researchers may wish to exercise caution in making claims about the “validity” of the noninvasive approach, which, at present, remains completely unproven, especially in renal disease.

ELDON D. LEHMANN
London, England, United Kingdom

Correspondence to Eldon D. Lehmann, M.B., B.S., B.Sc., Department of Imaging (MR Unit), National Heart and Lung Institute, Imperial College of Science Technology and Medicine, Royal Brompton Hospital, Sydney Street, London SW3 6NP, United Kingdom.
E-mail: info-www@2aida.org

REFERENCES

1. COVIC A, GOLDSMITH DJA, PANAGHIU L, *et al*: Analysis of the effect of hemodialysis on peripheral and central arterial pressure waveforms. *Kidney Int* 57:2634–2643, 2000
2. KARAMANOGU M, O’ROURKE MF, AVOLIO AP, KELLY RP: An analysis of the relationship between central aortic and peripheral upper limb pressure waves in man. *Eur Heart J* 14:160–167, 1993
3. O’ROURKE MF: Method for ascertaining the pressure pulse and related parameters in the ascending aorta from the contour of the pressure pulse in the peripheral arteries. *United States Patent No. 5,265,011*, 1993
4. FETICS B, NEVO E, CHEN C-H, KASS DA: Parametric model derivation of transfer function for noninvasive estimation of aortic pressure by radial tonometry. *IEEE Trans Biomed Eng* 46:698–706, 1999

Reply from the author

We thank Dr. Eldon Lehmann for his letter and the interest he has shown in our recent publication [1].

Over the last 2 years Dr. Lehmann, a noted expert in vascular imaging and methodology, with a long-standing interest in arterial compliance, has repeatedly censured other authors of other studies in which similar methods have been used [2–4].

Interested readers are urged to follow this correspondence trail across time, and several journals, the better to appreciate the background to these comments.

The thrust of his comments to us can be summarized as follows. First, is there any justification/supportive evidence for the use of a reverse generalized energy transfer function (GTF), as opposed to an individualized energy transfer function? Second, which parameters can safely be derived using a validated-GTF? Third, has there been any independent validation of the use of such GTFs with noninvasively calibrated brachial artery blood pressure? Finally, have the methodology and algorithms in use in the SphygmoCor device (PWV Medical, Inc., Sydney, Australia) been validated? Our answers (for brevity) are, “yes; aortic systolic blood pressure and augmentation index; yes, but not yet in the public domain; and partly.” We have reason to hope that later this year the answer to all of the questions will be “yes.” Until then, we concede that complete validation of this extremely interesting and potentially useful pulse wave analysis technique is (eagerly) awaited.

DAVID J. A. GOLDSMITH, for the authors
London, England, United Kingdom

Correspondence to David J.A. Goldsmith, M.A., F.R.C.P., Renal Unit, 4th Floor Thomas Guy House, Guy’s Hospital, St. Thomas’ Street, London, SE1 9RT England, United Kingdom.